



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Quantifying the association between physical activity and cardiovascular disease

Citation for published version:

Wahid, AA, Manek, N, Nichols, M, Kelly, P, Foster, C, Webster, P, Kaur, A, Smith, CF & Scarborough, P, 2015, 'Quantifying the association between physical activity and cardiovascular disease: A meta-analysis', *Cardiology in the young*, vol. 131, pp. 92-92.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cardiology in the young

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



QUANTIFYING THE ASSOCIATION BETWEEN PHYSICAL ACTIVITY AND CARDIOVASCULAR DISEASE: A META-ANALYSIS

A.A. Wahid¹, N. Manek², M. Nichols⁴, P. Kelly⁵, C. Foster³, P. Webster⁶, A. Kaur³, C. Friedemann Smith³, P. Scarborough³

1. University of Oxford, Oxon, UK

2. Watford General Hospital, Imperial College Healthcare NHS Trust, London, UK

3. The British Heart Foundation Centre on Population Approaches for Non-Communicable Disease Prevention, Department of Public Health, University of Oxford, UK

4. WHO Collaborating Centre for Obesity Prevention, Deakin University, Geelong, Australia

5. Sport, Physical Education and Health Sciences Department (SPEHS), University of Edinburgh, UK

6. University of Oxford Public Health Department, UK

Background and objective: The relationships between physical activity (PA) and cardiovascular disease (CVD) have predominantly been estimated using categorical measures of PA. In this systematic review and meta-analysis we derive a single continuous physical activity metric to directly compare the association between activity and CVD, both before and after adjustment for a measure of body weight.

Data sources: A systematic review was conducted through searching electronic databases such as MEDLINE and EMBASE for studies published between 1981 and 2014.

Study eligibility criteria and participants: Prospective cohort studies were included that measured PA levels where at least two of the following domains were measured: leisure, active travel and occupational activity. The relative risk needed to have been reported in healthy individuals and been adjusted for a measure of body weight. The PA exposure in each study was converted to MET hours per day. Various transformations were explored to parametrically describe the dose-response relationships, as well as a non-parametric categorical approach.

Results: A total of thirty-six studies were included in the analysis. An increase from inactive to achievement of recommended PA levels 150 minutes of moderate-intensity aerobic activity reduced the risk of CVD mortality by 23% and CVD incidence by 17% (RR 0.77 (0.71-0.84) and (RR 0.83 (0.77-0.89) respectively, after adjustment for body weight. Overall, there were a total of 3,439,874 participants, with 179,393 events occurring during an average follow up period of 12.3 years.

Conclusions: and implications: A single continuous metric for PA levels allowed us to directly compare the effect of physical activity on CVD incidence and mortality including myocardial infarct (MI), coronary heart disease (CHD) and heart failure. Effect sizes suggested that the greatest gain in health is associated with moving from inactive to small amounts of physical activity.